arXiv:math/0510279v3 [math.SP] 29 Jan 2008

FUNDAMENTAL TONE, CONCENTRATION OF DENSITY TO POINTS AND CONFORMAL DEGENERATION ON SURFACES

ALEXANDRE GIROUARD

Abstract. We study the effect of two types of degeneration of the Riemannian metric on the first eigenvalue of the Laplace operator on surfaces. In both cases we prove that the first eigenvalue of the round sphere is an optimal asymptotic upper bound. The first type of degeneration is concentration of the density to a point within a conformal class. The second is degeneration of the conformal class to the boundary of the moduli space on the torus and on the Klein bottle. In the latter, we follow the outline proposed by N. Nadirashvili in 1996.

1. Introduction

Given a Riemannian metric g on a closed surface Σ, let the spectrum of the Laplace operator Δ_g acting on smooth functions be the sequence

$$0 = \lambda_0(g) < \lambda_1(g) \leq \lambda_2(g) \leq \cdots \leq \lambda_k(g) \leq \cdots \nearrow \infty$$

where each eigenvalue is repeated according to its multiplicity. The first nonzero eigenvalue $\lambda_1(g)$ is called the *fundamental tone* of (Σ, g). Let $\mathcal{R}(\Sigma)$ be the space of Riemannian metrics on Σ with total area one. We are interested in the asymptotic behavior of the functional

$$\lambda_1 : \mathcal{R}(\Sigma) \rightarrow]0, \infty[$$

under two types of degeneration of sequences of metrics of area one $(g_n) \subset \mathcal{R}(\Sigma)$ derscribed below.

1.1. **Concentration to points.**

It is expected that a metric maximizing $\lambda_1 : \mathcal{R}(\Sigma) \rightarrow]0, \infty[$ has a lot of symmetries. On the sphere, the torus and the projective plane for example, the λ_1-maximizing metrics are the standard homogeneous ones. We consider the opposite situation where the distribution of mass of a sequence of metrics concentrate to a point, developping a δ-like singularity.

Date: February 2, 2008.

2000 *Mathematics Subject Classification.* Primary 35P, Secondary 58J.

Definition 1.1.1. A sequence $(g_n) \subset \mathcal{R}(\Sigma)$ is said to *concentrate to the point* $p \in \Sigma$ if for each neighborhood $\mathcal{O}$ of p

$$\lim_{n\to\infty} \int_{\mathcal{O}} dg_n = 1.$$

Question. Does concentration to a point impose any restriction to the asymptotic behavior of the eigenvalues of the Laplace operator Δ_{g_n} on the surface Σ?

Without any further constraints, the answer is no.

Proposition 1.1.2. *For any metric g_0, and any point $p \in \Sigma$ there exists a sequence (g_n) of pairwise isometric metrics concentrating to p. In particular the metrics (g_n) are isospectral.*

Under the additional assumption that the metrics g_n are conformally equivalent, we obtain an optimal asymptotic upper bound on the fundamental tone.

Theorem 1.1.3. *Let $[g] = \{\alpha g \mid \alpha \in C^\infty(\Sigma), \alpha > 0\}$ be a conformal class on a closed surface Σ.*

a) *For any sequence (g_n) of metrics of area one in the conformal class $[g]$, there holds*

$$\limsup_{n\to\infty} \lambda_1(g_n) \leq 8\pi.$$

b) *For any point $p \in \Sigma$, there exists a sequence (g_n) of metrics of area one in the conformal class $[g]$ concentrating to p and such that*

$$\lim_{n\to\infty} \lambda_1(g_n) = 8\pi.$$

Proposition 1.1.2 and Theorem 1.1.3 will be proved in section 5.

1.2. **Conformal degeneration.**

Given a conformal class $[g]$ on the torus T^2, define

$$\nu([g]) := \sup_{\tilde{g}\in\mathcal{R}(T^2)\cap[g]} \lambda_1(\tilde{g}).$$

This corresponds to the first conformal eigenvalue of Colbois and El Soufi [5]. Let

$$\mathcal{M} := \{a + ib \in \mathbb{C} \mid 0 \leq a \leq 1/2, a^2 + b^2 \geq 1, b > 0\}.$$

Any metric on T^2 is conformally equivalent to a flat torus $\mathbb{C}/\Gamma$ for some lattice Γ of $\mathbb{C}$ generated by $1 \in \mathbb{C}$ and $a + ib \in \mathcal{M}$. It follows that $\mathcal{M}$ is a natural representation of the moduli space $\mathcal{M}(T^2)$ of conformal classes on the torus (See Figure 1).



FIGURE 1. Moduli space of conformal structures on T^2

Definition 1.2.1. A sequence of metrics on the torus T^2 is *degenerate* if the corresponding sequence $(a_n+ib_n)\subset\mathcal{M}$ satisfies $\lim_{n\to\infty} b_n=\infty$.

Theorem 1.2.2. *If a sequence (g_n) of Riemannian metrics of area one on the torus is degenerate, then*

$$\lim_{n\to\infty}\nu([g_n])=8\pi.$$

In particular,

$$\limsup_{n\to\infty}\lambda_1(g_n)\le 8\pi.$$

The proof will be presented in section 3. We complete the outline proposed by Nadirashvili in [16]. In particular, a detailed version of a concentration lemma (Lemma 3.2.1) is used and a Dirichlet energy estimate on long cylinders is proposed (Lemma 3.6.1 and Lemma 2.1.1).

1.3. **Maximization of λ_1 on surfaces.**

One motivation for Theorem 1.2.2 is the role that it plays in λ_1-maximization on the torus. More generally, it is natural to ask which metric on a closed surface Σ, if any, maximizes the fundamental tone λ_1 in the space $\mathcal{R}(\Sigma)$ of Riemannian metrics of area one. It is known that

$$\Lambda(\Sigma):=\sup_{g\in\mathcal{R}(\Sigma)}\lambda_1(g)$$

is finite [10], [21], [19] for any closed surface Σ, but the explicit value of $\Lambda(\Sigma)$ has only been computed for few surfaces.

The first such result was obtained by Hersch [10] in 1970. He proved that the round metric g_{S^2} of area one is the unique maximum of λ_1 on

$\mathcal{R}(S^2)$. The proof rests on the fact that by Riemann's uniformization theorem any two metrics on the sphere are conformally equivalent.

In 1973, Berger [2] proved that among flat metrics on the torus, λ_1 is maximized by the flat equilateral metric g_{eq}, that is, the metric induced from the quotient of $\mathbb{C}$ by the lattice generated by 1 and $e^{i\pi/3}$(indicated by the $*$ in Figure 1). He conjectured that this metric is a global maximum of λ_1 over all Riemannian metrics of total area one. In 1996, Nadirashvili [16] proposed a method of proof.

Nadirashvili's approach. Start with a maximising sequence (g_n), i.e. such that $\lambda_1(g_n) \to \Lambda(T^2)$, and show that it admits a subsequence converging to a real analytic metric $\overline{g}$. Nadirashvili [16] has proved that a metric maximizing λ_1 on a surface Σ is also λ_1-minimal, which means that $(\Sigma, \overline{g})$ is minimally immersed in a round sphere by its first eigenfunctions. Montiel and Ros [15] have proved that for any surface different from the sphere (in particular for the torus!), the isometry group of a λ_1-minimal metric $\overline{g}$ coincide with its group of conformal transformations. See also [6]. Since the group of conformal transformations of a torus acts transitively, this implies that the curvature of $\overline{g}$ is constant and therefore zero by Gauss-Bonnet theorem. The above result of Berger [2] completes the proof.

The first step in showing that (g_n) admits a convergent subsequence is to prove that the associated sequence $([g_n])$ of conformal classes admits a converging subsequence. Since $\lambda_1(g_{eq}) = 8\pi^2/\sqrt{3} > 8\pi$, Theorem 1.2.2 implies that the corresponding sequence (a_n+ib_n) is bounded and must therefore admits a convergent subsequence.

Remark 1.3.1.

- Explicit λ_1-maximal metrics are also known for the projective space [19] and the Klein bottle [13], [7]. There is a conjecture for surfaces of genus two [12].
- The existence of analytic λ_1-maximal metrics has recently been used in [13] and [7].

1.4. **Conformal degeneration on the Klein bottle.**

Define two affine transformations t_b and τ of $\mathbb{C}$ by

$$t_b(x+iy) = x + i(y+b),$$
$$\tau(x+iy) = x + \pi - iy.$$

Let G_b be the group of transformations generated by t_b and τ.

Lemma 1.4.1. *Any Riemannian metric g on the Klein bottle $\mathbb{K}$ is conformally equivalent to one of the standard flat models*

$$K_b := \mathbb{C}/G_b.$$

In other words, there exist a smooth function $\alpha : K_b \to]0,\infty[$ such that $(\mathbb{K}, g)$ is isometric to $(K_b, \alpha(dx^2 + dy^2))$.

It follows that the moduli space of conformal classes on the Klein bottle is identified with the set of positive real numbers.



Theorem 1.4.2. *Let $(g_n) \subset \mathcal{R}(\mathbb{K})$ be a sequence of metrics on the Klein bottle.*

1. *If* $\lim_{n\to\infty} b_n = 0$, *then* $\limsup_{n\to\infty} \lambda_1(g_n) = 8\pi$.
2. *If* $\lim_{n\to\infty} b_n = \infty$, *then* $\limsup_{n\to\infty} \lambda_1(g_n) \leq 12\pi$.

The proof will be presented in section 4, we follow the outline proposed by Nadirashvili in [16] and in a private communication. The first case is very similar to the corresponding result for the torus (Theorem 1.2.2). The second case uses the fact that the standard metric on $\mathbb{R}P^2$ is minimally embedded in S^4 by its first eigenfunctions to obtain an estimate on the Dirichlet energy of a test function via a theorem of Li and Yau [19] on conformal area of minimal surfaces.

1.5. **Friedlander and Nadirashvili invariant.** For a closed manifold M of dimension at least 3, Colbois and Dodziuk [4] proved that the first eigenvalue is unbounded on the set of Riemannian metrics of area one, that is $\Lambda(M) = +\infty$. On the other hand, it is known that the supremum $\nu(C)$ of λ_1 restricted to metrics of area one in any fixed conformal class C is finite [8]. Friedlander and Nadirashvili [17] introduced the differential invariant

$$I(M) := \inf\left\{\nu(C) \;\middle|\; C \text{ is a conformal class on } M\right\}$$

and proved that it satisfies

$$I(M) \geq \lambda_1(S^n, g_{S^n})$$

where g_{S^n} is the round metric of area one on the sphere S^n. It would be interesting to know if this invariant distinguishes nonequivalent differential structures. In fact, it is very difficult to compute $I(\Sigma)$ explicitely, even for surfaces. Since any two metrics on S^2 are conformally equivalent, it is obvious that $I(S^2) = 8\pi$. For similar reasons, $I(\mathbb{R}P^2) = 12\pi$. The invariant for the torus and for the Klein bottle are obtained as corollaries to Theorem 1.2.2 and Theorem 1.4.2.

Corollary 1.5.1. *The Friedlander-Nadirashvili invariants the torus and the Klein bottle are* 8π*.*

Let us also mention the following conjecture.

Conjecture 1.5.2 (Friedlander-Nadirashvili)**.** *For any closed surface* Σ *other than the projective plane* $\mathbb{R}P^2$*,* $I(\Sigma) = 8\pi$*.*

2. Analytic background

Let Σ be a closed surface. In dimension two the Dirichlet energy of a function $u \in C^\infty(\Sigma)$

$$D(u) = \int_\Sigma |\nabla_g u|^2 \, dg$$

is invariant under conformal diffeomorphisms. In order to estimate the first eigenvalue of the Laplace operator Δ_g, the following variational characterization will be used:

$$\lambda_1(g) = \inf \left\{ \frac{D(u)}{\int_\Sigma u^2 \, dg} \;\middle|\; u \in C^\infty(\Sigma), u \neq 0, \int_\Sigma u \, dg = 0 \right\}. \tag{2.0.1}$$

2.1. Dirichlet energy estimate on thin cylinders.

The main technical tool that we use is an estimate on the Dirichlet energy of harmonic functions on long cylinders in terms of their restrictions to the boundary circles.

Lemma 2.1.1. *Let* $\Omega = (0, L) \times S^1$ *with* $S^1 = \mathbb{R}/2\pi\mathbb{Z}$*. Consider* $f \in C^\infty(\overline{\Omega})$ *such that* $f(0, \theta) = 0$ *and* $|f(L, \theta)| \leq 1$*. Let* $u(\theta) = f(L, \theta)$*. If* f *is harmonic, then*

$$\int_\Omega |\nabla f|^2 \leq \frac{2\pi}{L} + \coth(L) \int_{\theta=0}^{2\pi} |u'(\theta)|^2 d\theta.$$

The idea is to express the Fourier series of the function f in terms of the Fourier series of u. This is similar to Hurwitz's proof of the isoperimetric inequality [11].

Proof. Let the Fourier representation of u be

$$u(\theta) = K + \sum_{k=1}^{\infty} \big(a_k \cos(k\theta) + b_k \sin(k\theta)\big).$$

Direct computation shows that f admits the following representation:

$$f(x,\theta) = \frac{Kx}{L} + \sum_{k=1}^{\infty} \frac{\sinh(kx)}{\sinh(kL)} \Big(a_k \cos(k\theta) + b_k \sin(k\theta)\Big)$$

and integration by parts leads to

$$\begin{aligned}
\int_\Omega |\nabla f|^2 &= -\int_\Omega f \Delta f + \int_{\partial\Omega} f \frac{\partial f}{\partial \nu} \\
&= \int_{\theta=0}^{2\pi} u(\theta) \partial_x f(x,\theta) d\theta \Big|_{x=L} \\
&= \int_{\theta=0}^{2\pi} \Big(K + \sum_{k=1}^{\infty} a_k \cos(k\theta) + b_k \sin(k\theta)\Big) \times \\
&\qquad \Big(\frac{K}{L} + \sum_{k=1}^{\infty} k \coth(kL) \left(a_k \cos(k\theta) + b_k \sin(k\theta)\right)\Big) d\theta \\
&= 2\pi \frac{K^2}{L} + \pi \sum_{k=1}^{\infty} k \coth(kL) (a_k^2 + b_k^2)
\end{aligned}$$

For $x > 0$,

$$\frac{d}{dx} \coth(x) = -\frac{4}{(e^x - e^{-x})^2} < 0$$

so that for each $k \geq 1$,

$$\coth(kL) \leq \coth(L).$$

It follows that

$$\begin{aligned}
\int_\Omega |\nabla f|^2 &\leq 2\pi \frac{K^2}{L} + \pi \coth(L) \sum_{k=1}^{\infty} k (a_k^2 + b_k^2) \\
&\leq 2\pi \frac{K^2}{L} + \pi \coth(L) \sum_{k=1}^{\infty} k^2 (a_k^2 + b_k^2) \\
&= 2\pi \frac{K^2}{L} + \coth(L) \int_{\theta=0}^{2\pi} |u'(\theta)|^2 d\theta
\end{aligned}$$

where $|K| = \frac{1}{2\pi} |\int_{\theta=0}^{2\pi} u(\theta) d\theta| \leq 1$ since $u(\theta) \in [-1, 1]$. $\square$

A simple conformal change of coordinates is used to extend Lemma 2.1.1 to the situation where the boundary circle has arbitrary length.

Corollary 2.1.2. *Suppose the hypothesis of Lemma 2.1.1 holds with the circle $\mathbb{R}/2\pi\mathbb{Z}$ replaced by $\mathbb{R}/r\mathbb{Z}$. For any $L > 0$*

$$\int_\Omega |\nabla f|^2 \leq \frac{r}{L} + \frac{r}{2\pi}\coth(\frac{2\pi L}{r}) \int_{x=0}^{r} |u'(x)|^2 dx. \tag{2.1.1}$$

2.2. **Conformal renormalization of centers of mass.** It is possible to conformally move any nonsingular distribution of mass on the sphere $S^n \subset \mathbb{R}^{n+1}$ in such a way that its center of mass becomes the origin of $\mathbb{R}^{n+1}$.

Lemma 2.2.1 (Hersch Lemma)**.** *Let μ be a measure on the sphere S^n. If the support of μ is not a point, then there exists a conformal transformation τ of the sphere S^n such that*

$$\int_{S^n} \pi \circ \tau \; d\mu = 0 \tag{2.2.1}$$

where $S^n \overset{\pi}{\hookrightarrow} \mathbb{R}^{n+1}$ is the standard embedding.

This lemma was obtained by Hersch [10] in 1970, see also [20]. It is proved using a topological argument similar to the proof of Brouwer fixed point theorem.

Corollary 2.2.2. *Let μ be a measure on a surface Σ. Consider an embedding $\phi : \Sigma \longrightarrow S^n$. If the support of μ is not a point, then there exists a conformal transformation τ of S^n such that*

$$\int_\Sigma \pi \circ \tau \circ \phi \, d\mu = 0.$$

Proof. The result follows from application of Hersch Lemma to the push-forward measure $\phi_*\mu$ since $\int_{S^n} f \, d(\phi_*\mu) = \int_\Sigma f \circ \phi \, d\mu$ for any smooth function f. □

3. CONFORMAL DEGENERATION ON THE TORUS

The goal of this section is to prove Theorem 1.2.2.

3.1. **Moduli space of tori.** For any $b > 0$, let

$$T_b := \Big\{[x+iy] \in \mathbb{C}/\mathbb{Z} \;\Big|\; -b/2 < y < b/2\Big\}$$

be a cylinder of length b. Given $a + ib \in \mathcal{M}$, let $\Gamma_{a,b}$ be the lattice of $\mathbb{C}$ generated by 1 and $a + ib$. Consider the group $G_{a,b}$ of transformations of T_∞ generated by

$$[x+iy] \mapsto [x+a+i(y+b)].$$

The cylinder T_b is a fundamental domain of this action and the torus $\mathbb{C}/\Gamma_{a,b}$ can also be obtained as $T_\infty/G_{a,b}$.

3.2. **Concentration on thin cylinders.** In order to make notation less cumbersome, consider a sequence (g_n) of metrics such that $b_n = n$. The first eigenvalue of the flat torus corresponding to g_n,

$$\lambda_1(T_n/G_{a_n,n}) = \frac{4\pi^2}{n^2}$$

tends to zero with n going to infinity [1]. Imposing a uniform positive lower bound $\lambda_1(g_n) \geq K > 0$ on the first eigenvalues for g_n should therefore imply that g_n is "far from being flat". The next lemma makes this intuitive idea precise by showing that the Riemannian measures dg_n concentrate on relatively thin cylindrical parts in (T^2, g_n).

Lemma 3.2.1. *If there exists $K > 0$ such that*

$$\liminf_{n\to\infty} \lambda_1(g_n) \geq K,$$

then for any $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for any $n \geq N$,

$$\max\{\int_{T_{3n/4}} dg_n, \int_{T_n \setminus T_{n/4}} dg_n\} \geq 1 - \epsilon.$$

Let A_n^ϵ be the maximizing cylinder: either $T_{3n/4}$ or $T_n \setminus T_{n/4}$. This lemma says that most of the mass (i.e. $1-\epsilon$) is concentrated on a cylinder whose lenth is $3/4$ of the total length. Without loss of generality, we will suppose $A_n^\epsilon = T_{3n/4}$.

Proof. The function

$$\gamma_n([x+iy]) = \cos(\frac{2\pi y}{n})$$

is a first eigenfunction on the flat torus $T_n/G_{a_n,n}$ corresponding to g_n. Let



FIGURE 2. Concentration

$$c_n = \int_{T_n} \gamma_n(x+iy)\, dg_n$$

and define $h_n : T_n \to \mathbb{R}$ by

$$h_n = \gamma_n - c_n,$$

which satisfies

$$\int_{T_n} h_n \, dg_n = 0.$$

Given $\delta > 0$, let $P_n^\delta = h_n^{-1}([-\delta, \delta])$ and $Q_n^\delta = T_n \setminus P_n^\delta$. Since $\delta^2 \leq h_n^2$ on Q_n^δ,

$$\delta^2 \int_{Q_n^\delta} dg_n \leq \int_{Q_n^\delta} h_n^2 \, dg_n \leq \int_{T_n} h_n^2 \, dg_n$$

Using the variational characterization of $\lambda_1(g_n)$ and the conformal invariance of the Dirichlet energy,

$$\delta^2 \int_{Q_n^\delta} dg_n \leq \frac{\int_{T_n} |\nabla h_n|^2 \, dg_n}{\lambda_1(g_n)} \leq \frac{\int_{T_n} |\nabla \gamma_n|^2 \, dg_n}{K} = \frac{4\pi^2}{Kn^2} \to 0.$$

So that, for each $\delta > 0$,

$$\lim_{n\to\infty} \int_{Q_n^\delta} dg_n = 0$$

and

$$\lim_{n\to\infty} \int_{P_n^\delta} dg_n = 1.$$

Observe that

$$P_n^\delta = \left\{ [x+iy] \;\middle|\; c_n - \delta \leq cos(2\pi y/n) \leq c_n + \delta \right\}.$$

For $\delta > 0$ small enough, P_n^δ is the union of two intervals around $\pm \arccos(c_n)$ whose lengths are at most $\frac{n}{10}$. The set P_n^δ will either be included in $T_{3n/4}$ or in $T_n \setminus T_{n/4}$ (see Figure 2). $\square$

3.3. **Transplantation to the sphere.** Let $\sigma : \mathbb{C} \to S^2$ be the stereographic parametrization of the sphere by its equatorial plane

$$\sigma(u+iv) = \frac{1}{1+u^2+v^2}(2u, 2v, u^2+v^2-1) \tag{3.3.1}$$

and define the conformal equivalence $\phi : \mathbb{C}/\mathbb{Z} \to S^2 \setminus \{\text{poles}\}$ by

$$\phi([z]) = \sigma(e^{-2\pi i z}).$$

3.4. **Renormalization of the centers of mass.** It follows from Corollary 2.2.2 that there exists conformal transformations τ_n such that

$$\int_{A_n^\epsilon} \pi \circ \tau_n \circ \phi \, dg_n = 0 \tag{3.4.1}$$

where $S^2 \overset{\pi}{\hookrightarrow} \mathbb{R}^3$ is the standard embedding. For each $n \in \mathbb{N}$, since $\pi_1^2 + \pi_2^2 + \pi_3^2 = 1$ on S^2, there exists an indice $i = i(n) \in \{1, 2, 3\}$ such that the function

$$u_n = \pi_i \circ \tau_n \circ \phi \tag{3.4.2}$$

satisfies

$$\int_{A_n^\epsilon} u_n^2 \, dg_n \geq \frac{1}{3} \int_{A_n^\epsilon} dg_n \geq \frac{1-\epsilon}{3}. \tag{3.4.3}$$

3.5. **Test functions.** The function u_n will be extended and perturbed to a function f_n defined on T_n and admissible for the variational characterization (2.0.1) of $\lambda_1(g_n)$.

Let $I_n^- = [-7n/16, -6n/16]$ and $I_n^+ = [6n/16, 7n/16]$.



Given $\alpha_n^- \in I_n^-$ and $\alpha_n^+ \in I_n^+$, define cylinders

$$B(\alpha_n^-) = \left\{ [x+iy] \in T_n \;\middle|\; y \leq \alpha_n^- \right\},$$
$$B(\alpha_n^+) = \left\{ [x+iy] \in T_n \;\middle|\; \alpha_n^+ \leq y \right\}.$$



Their lengths are at least $n/16$. Define $w_n : T_n \to \mathbb{R}$ by the following differential problem:

$$\begin{cases} \Delta w_n = 0 & \text{on } B(\alpha_n^-) \cup B(\alpha_n^+), \\ w_n = 0 & \text{on } \partial T_n = \left\{ [x+iy] \in \mathbb{C}/\mathbb{Z} \;\middle|\; |y| = n/2 \right\}, \\ w_n = u_n & \text{on } T_n \setminus \left(B(\alpha_n^-) \cup B(\alpha_n^+) \right). \end{cases}$$

Since the continuous function w_n is piecewise smooth and satisfies $w_n = 0$ on the boundary of T_n, it is compatible with the identification of the boundary and induces a piecewise smooth function on the torus.

Let $\delta_n = \int_{T_n} w_n\, dg_n$. Since $\int_{A_n^\epsilon} w_n\, dg_n = 0$, it follows from concentration of the measures dg_n on A_n^ϵ and from the maximum principle that

$$|\delta_n| = \left|\int_{T_n\setminus A_n^\epsilon} w_n\, dg_n\right| \leq \max_{x\in T_n\setminus A_n^\epsilon} |w_n(x)| \int_{T_n\setminus A_n^\epsilon} dg_n \leq \varepsilon.$$

This means that w_n is almost admissible for the variational characterization of $\lambda_1(g_n)$. Define $f_n : T_n \to \mathbb{R}$ by

$$f_n = w_n - \delta_n \tag{3.5.1}$$

so that $\int_{T_n} f_n\, dg_n = 0$. From (3.4.1) and (3.4.3) it follows that

$$\begin{aligned} \int_{T_n} f_n^2\, dg_n &\geq \int_{A_n^\epsilon} (u_n - \delta_n)^2\, dg_n \\ &= \int_{A_n^\epsilon} u_n^2\, dg_n + \delta_n^2 \int_{A_n^\epsilon} dg_n \geq \frac{\mathcal{A}(A_n^\epsilon, g_n)}{3} \geq \frac{1-\epsilon}{3} \end{aligned} \tag{3.5.2}$$

Using the variational characterization (2.0.1) of $\lambda_1(g_n)$ and conformal invariance of the Dirichlet energy leads to

$$\begin{aligned} \lambda_1(g_n) &\leq \frac{3\int_{T_n} |\nabla f_n|^2\, dg_n}{1-\varepsilon} \\ &= \frac{3}{1-\varepsilon}\left(\int_{T_n\setminus(B(\alpha_n^-)\cup B(\alpha_n^+))} |\nabla w_n|^2\, dg_n \right. \\ &\qquad\qquad \left. + \int_{B(\alpha_n^-)\cup B(\alpha_n^+)} |\nabla w_n|^2\, dg_n\right) \\ &\leq \frac{3}{1-\varepsilon}\left(\int_{S^2} |\nabla \pi_i|^2\, dg_{S^2} + \int_{B(\alpha_n^-)\cup B(\alpha_n^+)} |\nabla w_n|^2\, dg_n\right) \\ &= \frac{8\pi}{1-\varepsilon} + \frac{3}{1-\varepsilon}\int_{B(\alpha_n^-)\cup B(\alpha_n^+)} |\nabla w_n|^2\, dg_n. \end{aligned} \tag{3.5.3}$$

3.6. **Energy estimate.** On a long flat cylinder like $B(\alpha_n^\pm)$, the Dirichlet energy of a harmonic function is controled by the Dirichlet energy of its restriction to the boundary circles. Corollary 2.1.2 is applied to the function $f = w_n$ on the cylinders $\Omega = B(\alpha_n^\pm)$, their lengths are at least $n/16$. For any x, $u(x) := f(x, \alpha_n^\pm) \in [-1, 1]$ since it is a coordinate function on the sphere.

The next lemma shows that the numbers $\alpha_n^\pm \in I_n^\pm$ can be chosen to make the Dirichlet energy of u_n on the boundary of $B(\alpha_n^\pm)$ small.

Lemma 3.6.1. *There exist $\alpha_n^- \in I_n^-$ and $\alpha_n^+ \in I_n^+$ such that*

$$\int_{x=0}^1 |\partial_x u_n(x+i\alpha_n^\pm)|^2 dx \leq \frac{128\pi}{3n}.$$

Proof. Let $E_n^\pm = \left\{[x+iy] \;\middle|\; y \in I_n^\pm\right\}$. Since the width of $I_n^\pm$ is at least $n/16$, the mean-value theorem implies

$$\iint_{E_n^\pm} |\nabla u_n|^2 \; dxdy \geq \frac{n}{16} \min\left\{\int_{x=0}^1 |\nabla u_n(x+i\alpha_n^\pm)|^2 dx \;\middle|\; \alpha_n^\pm \in I_n^\pm\right\}$$

In particular, there exists $\alpha_n^\pm \in I_n^\pm$ such that

$$\int_{x=0}^1 |\partial_x u_n(x+i\alpha_n^\pm)|^2 dx \leq \frac{16}{n}\iint_{E_n^\pm} |\nabla u_n|^2 \; dxdy.$$

By conformal invariance of the Dirichlet energy, this is bounded above by

$$\frac{16}{n}\int_{\tau_n\circ\phi(E_n^\pm)} |\nabla \pi_i|^2 \, dg_{S^2} \leq \frac{16}{n}\int_{S^2} |\nabla \pi_i|^2 \, dg_{S^2} = \frac{16}{n}.\frac{8\pi}{3} = \frac{128\pi}{3n}.$$

□

Proof of Theorem 1.2.2. Let f_n be the function given by (3.5.1). Using the estimate on boundary derivative (Lemma 3.6.1), the Dirichlet energy estimate on cylinders (Lemma 2.1.1) implies

$$\lim_{n\to\infty} \int_{B(\alpha_n^\pm)} |\nabla f_n|^2 \, dg_n = 0. \tag{3.6.1}$$

Using inequality (3.5.3), obtained by the variational characterization of $\lambda_1(g_n)$, it follows that $\limsup_n \lambda_1(g_n) \leq \frac{8\pi}{1-\epsilon}$. Since $\epsilon > 0$ is arbitrary, it follows that

$$\limsup_{n\to\infty} \lambda_1(g_n) \leq 8\pi.$$

Finally, the lower bound follows from the result of Friedlander and Nadirashvili [17] stated in the introduction: for any conformal class C on a closed surface, $\nu(C) \geq 8\pi$. □

4. Conformal degeneration on the Klein bottle

The goal of this section is to prove Theorem 1.4.2. Let $S^k(r)$ be the k-dimentional sphere of radius r with its standard metric $g_{S^k(r)}$ and $\mathbb{R}P^2(r)$ be the associated projective plane with standard metric $g_{\mathbb{R}P^2(r)}$. Recall from the introduction that any Klein bottle is conformally equivalent to a unique $K_b = \mathbb{C}/G_b$ where G_b is the group of transformations of $\mathbb{C}$ generated by $t_b(x+iy) = x + i(y+b)$ and

$\tau(x+iy) = x + \pi - iy$. The rectangle $[0,\pi] \times [-b/2, b/2]$ is a fundamental domain for the action of G_b on $\mathbb{C}$. Reversing and identifying the opposite vertical sides this rectangle, we obtain a Möbius strip

$$M_b = \big([0,\pi] \times [-b/2, b/2]\big)/\tau.$$

4.1. **Transplantation to the sphere S^4 via projective space.** In this paragraph we exhibit a conformal embedding of the infinite Möbius strip M_∞ in the sphere S^4. We start with a lemma which will be used to embed Möbius strip conformally in $\mathbb{R}P^2$.

Lemma 4.1.1. *The conformal application $\phi : \mathbb{C} \to S^2 \subset \mathbb{R}^3$ defined by*

$$\phi(x+iy) = \frac{1}{e^{2y}+1}(2e^y\cos(x), 2e^y\sin(x), e^{2y}-1)$$

satisfies

$$\phi(x+2\pi+iy) = \phi(x+iy),$$
$$\phi(\tau(x+iy)) = -\phi(x+iy).$$

It induces a conformal equivalence $\phi : M_\infty \to \mathbb{R}P^2 \setminus \{[0:0:1]\}$.

The Veronese map v is a well known minimal isometric embedding of $\mathbb{R}P^2(\sqrt{3})$ in the sphere S^4 by its first eigenfunctions. This means that the components of v are eigenfunctions for $\lambda_1(g_{\mathbb{R}P^2(\sqrt{3})}) = 2$. For details, see [1] and [14].

It follows that the composition $v \circ \phi$ is a conformal embedding of the Möbius strip M_∞ in S^4.

4.2. **Concentration on Möbius strips.** Without loss of generality, consider a sequence $b_n = n$.

Lemma 4.2.1. *If there exists $K > 0$ such that*

$$\liminf_{n\to\infty} \lambda_1(g_n) \geq K,$$

then for any $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that for any $n \geq N$,

$$\max\Big\{\int_{M_{3n/4}} dg_n, \int_{K_n \setminus K_{3n/4}} dg_n\Big\} \geq 1 - \epsilon.$$

Proof. Since the function

$$u_n([x+iy]) = \cos(\frac{2\pi y}{n})$$

used in the proof of Lemma 3.2.1 is even, it induces a first eigenfunction on the flat Klein bottle K_n. The cylinders constructed in this proof are compatible with the identification on the Möbius strip, also because the function u_n is even. □

Notation: Without loss of generality, we suppose the maximum is reached by $A_n^\epsilon := M_{3n/4}$. See Lemma 3.2.1 for details.

4.3. **Renormalization of the centers of mass.** It follows from Corollary 2.2.2 that there exists conformal transformations τ_n of S^4 such that

$$\int_{A_n^\epsilon} \pi \circ \tau_n \circ v \circ \phi \, dg_n = 0$$

where $\pi : S^4 \hookrightarrow \mathbb{R}^5$ is the standard embedding. For $1 \leq i \leq 5$, let

$$u_n^i = \pi_i \circ \tau_n \circ v \circ \phi. \tag{4.3.1}$$

4.4. **Test functions.** The numbers

$$\alpha_n \in I_n := [\frac{6n}{16}, \frac{7n}{16}],$$

will be chosen later (see Lemma 3.6.1). For each $1 \leq i \leq 5$, define $w_n^i : M_n \to \mathbb{R}$ by the following differential problem:

$$\begin{cases} \Delta w_n^i = 0 & \text{on } M_n \setminus M_{\alpha_n}, \\ w_n^i = 0 & \text{on } \partial M_n = \left\{ [(x,y)] \in M_n \;\middle|\; |y| = n/2 \right\}, \\ w_n^i = u_n^i & \text{on } M_{\alpha_n} \end{cases}$$

where u_n^i is defined by (4.3.1). Since the continuous function w_n^i is piecewise smooth and satisfies $w_n^i = 0$ on the boundary of M_n, it is compatible with the identification of the boundary and induces a piecewise smooth function on the Klein bottle K_n.

Since $\int_{A_n^\epsilon} w_n^i \, dg_n = 0$, it follows from concentration of the measures dg_n on A_n^ϵ and from the maximum principle that $\delta_n^i := \int_{M_n} w_n^i \, dg_n$ satisfies $|\delta_n^i| \leq \epsilon$. This means that w_n^i is almost admissible for the variational characterization of $\lambda_1(g_n)$. Thus, it is natural to define $f_n^i : M_n \to \mathbb{R}$ by

$$f_n^i = w_n^i - \delta_n^i \tag{4.4.1}$$

so that for each i

$$\int_{M_n} f_n^i \, dg_n = 0$$

and similarly to inequality (3.5.2)

$$\sum_{i=1}^5 \int_{M_n} (f_n^i)^2 \, dg_n \geq \int_{A_n^\epsilon} dg_n \geq 1 - \epsilon. \tag{4.4.2}$$

It follows from the variational characterization of $\lambda_1(g_n)$ and from inequality (4.4.2) that

$$\lambda_1(g_n)\,(1-\epsilon) \le \lambda_1(g_n)\left(\sum_{i=1}^{5}\int_{M_n}(f_n^i)^2\,dg_n\right) \le \sum_{i=1}^{5}\int_{M_n}|\nabla f_n^i|^2\,dg_n$$

$$= \sum_{i=1}^{5}\left(\int_{M_{\alpha_n}}|\nabla w_n^i|^2\,dg_n + \int_{M_n\setminus M_{\alpha_n}}|\nabla w_n^i|^2\,dg_n\right). \tag{4.4.3}$$

4.5. **Energy estimate.**

First step: Bounding $\sum_{i=1}^5\int_{M_{\alpha_n}}|\nabla w_n^i|^2\,dg_n$.

Recall that the Veronese embedding $v:\mathbb{R}P^2(\sqrt{3})\to S^4$ is isometric and minimal. On $\Sigma_n := \tau_n\circ v(\mathbb{R}P^2(\sqrt{3}))$ we consider the metric induced by g_{S^4}. Proposition 1 of [19] says that if a compact surface is minimally immersed in a sphere, then its area cannot be increased by conformal transformations of the sphere. In our particular case this leads to the following proposition.

Proposition 4.5.1. *For each* $n\in\mathbb{N}$,

$$\int_{\Sigma_n} dg_{S^4} \le \textit{Area of}\,\mathbb{R}P^2(\sqrt{3}) = 6\pi.$$

It follows from conformal invariance of the Dirichlet energy that

$$\sum_{i=1}^{5}\int_{M_{\alpha_n}}|\nabla w_n^i|^2\,dg_n \le \sum_{i=1}^{5}\int_{\Sigma_n}|\nabla\pi_i|^2\,dg_{S^4}$$

It is proved on page 146 of [20] that Σ_n being isometrically immersed in S^4 implies the point-wise identity $\sum_{i=1}^5|\nabla\pi_i|^2 = 2$. Whence

$$\sum_{i=1}^{5}\int_{M_{\alpha_n}}|\nabla w_n^i|^2\,dg_n \le 2\int_{\Sigma_n}dg_{S^4} \le 12\pi. \tag{4.5.1}$$

Second step: Bounding $\sum_{i=1}^5\int_{M_n\setminus M_{\alpha_n}}|\nabla w_n^i|^2\,dg_n$.

The function w_n is harmonic on the set $M_n\setminus M_{\alpha_n}$. This is a cylinder of length $L_n := n-\alpha_n \ge 9/16n$ and of width 2π. The next lemma shows that the Dirichlet energy of w_n on the boundary of these cylinders can be controlled by appropriate choice of α_n.

Lemma 4.5.2. *The number* $\alpha_n \in I_n = [6n/16, 7n/16]$ *can be chosen such that*

$$\sum_{i=1}^{5} \int_{x=0}^{\pi} |\partial_x w_n^i(x \pm i\alpha_n)|^2 dx \leq 192\pi/n.$$

Proof. We argue as in Lemma 3.6.1. Let $E_n^{\pm} = \{[x+iy] \mid y \in I_n\}$. It follows from the mean-value theorem that

$$\frac{n}{16} \min_{\alpha_n^{\pm} \in I_n^{\pm}} \sum_{i=1}^{5} \int_{x=0}^{2\pi} |\nabla w_n^i(x, \alpha_n^{\pm})|^2 dx \leq \sum_{i=1}^{5} \iint_{E_n^{\pm}} |\nabla w_n^i(x+iy)|^2 \, dxdy \\ \leq 12\pi.$$

☐

Since $M_n \setminus M_{\alpha_n}$ has length L_n at least $n/16$ and is of width 2π, Lemma 2.1.1 implies

$$\sum_{i=1}^{5} \int_{M_n \setminus M_{\alpha_n}} |\nabla w_n^i|^2 \, dg_n \\ \leq \sum_{i=1}^{5} \Big(\frac{2\pi}{L_n} + \coth(L_n) \int_{x=0}^{2\pi} |\partial_x w_n^i(x \pm i\alpha_n)|^2 \, dx \Big) \\ \leq \frac{\pi}{n} \left(160 + 5 \times 192 \coth(\frac{n}{16}) \right). \tag{4.5.2}$$

Proof of Theorem 1.4.2. Substitution of inequality (4.5.1) and inequality (4.5.2) in the variational characterization (4.4.3) leads to

$$\limsup_{n\to\infty} \lambda_1(g_n)(1-\epsilon) \leq 12\pi + \limsup_{n\to\infty} \frac{\pi}{n} (160 + 960 \coth(n/16)) = 12\pi.$$

Since $\epsilon > 0$ is arbitrary, this completes the proof of Theorem 1.4.2. ☐

5. Concentration to points

The main goal of this section is to prove Theorem 1.1.3. We start by proving that concentration to a point has no influence on spectrum.

Proof of Proposition 1.1.2. There exists a sequence of diffeomorphisms ϕ_n such that $\lim_{n\to\infty} \phi_n(x) = p$ dg_0-almost everywhere. Indeed, let $f : M \to \mathbb{R}$ be any Morse function having p as its unique local minimum and consider $\phi \mathbb{R} \times M \to M$ be its negative gradient flow with respect to g_0. Since the stable manifolds of any critical point other than p are of codimension strictly greater than 1, they are dg_0-negligible, hence the sequence ϕ_n satisfies the required property and $g_n = \phi_n^* g_0$ concentrates to p. ☐

We know proceed with the proof of Theorem 1.1.3.

5.1. **Construction of a neighborhood system.** Let $(g_n) \subset \mathcal{R}(\Sigma)$ be a sequence of metrics concentrating to $p \in \Sigma$. Let $\mathbb{D}$ be the unit open disk in $\mathbb{C}$. Let $\eta : D \to \mathbb{D}$ a conformal chart arround p such that $\eta(p) = 0$. Observe that since the metrics g_n are all in the same conformal class, the same chart η will be conformal for each of them.

Lemma 5.1.1. *There exists a conformal equivalence*

$$\psi : D \setminus \{p\} \to (0, \infty) \times S^1$$

and a family $\mathcal{U}_n \subset D$ *of neighborhood of* p *such that*

$$\lim_{n\to\infty} \int_{\mathcal{U}_n} dg_n = 1$$

and

$$\psi(D \setminus \mathcal{U}_n) = (0, L_n) \times S^1$$

with $L_n \to \infty$ *and* $\psi(\partial D) = \{0\} \times S^1$.

Proof. For $0 \leq \epsilon \leq 1$, let $\mathcal{U}(\epsilon) = \eta^{-1} B(0, \epsilon)$. Define ϵ_n by

$$\int_{\mathcal{U}(\epsilon_n)} dg_n = 1 - \epsilon_n.$$

It follows from concentration that $\lim_{n\to\infty} \epsilon_n = 0$ so that

$$\lim_{n\to\infty} \int_{\mathcal{U}(\epsilon_n)} dg_n = 1$$

Define the conformal equivalence $\alpha : \mathbb{R} \times S^1 \to \mathbb{C}^*$ by

$$\alpha(x, y) = e^{-x}\big(\cos(y), \sin(y)\big).$$

The composition $\psi = \alpha^{-1} \circ \eta$ has the required property since

$$\psi(D \setminus \mathcal{U}(\epsilon_n)) = \big(0, -\ln(\epsilon_n)\big) \times S^1.$$

□

5.2. **Renormalization of the centers of mass.** Recall that σ, as defined in (3.3.1), is the stereographic parameterization of the sphere S^2 by its equatorial plane. Let $H \subset S^2$ be the southern hemisphere of S^2. The map

$$\phi = \sigma \circ \eta : D \to H$$

is a conformal equivalence such that $\phi(p)$ is the south pole. It follows from Corollary 2.2.2 that there exist conformal transformations τ_n of the sphere such that

$$\int_{\mathcal{U}_n} \pi \circ \tau_n \circ \phi \, dg_n = 0$$

where $\pi : S^2 \hookrightarrow \mathbb{R}^3$ is the standard embedding. For each $n \in \mathbb{N}$, since $\pi_1^2 + \pi_2^2 + \pi_3^2 = 1$ on S^2, there exists an indice $i = i(n) \in \{1,2,3\}$ such that the function $u_n = \pi_i \circ \tau_n \circ \phi$ satisfies

$$\int_{\mathcal{U}_n} u_n^2 \, dg_n \geq \frac{1}{3} \int_{\mathcal{U}_n} dg_n.$$

5.3. **Test functions.** Consider $\alpha_n \in [L_n/2, L_n]$ to be chosen later. Define $w_n : \Sigma \to \mathbb{R}$ as the unique solution of

$$\begin{cases} w_n = 0 & \text{on } \Sigma \setminus D \\ w_n = \pi_i \circ \tau_n \circ \phi & \text{on } \mathcal{U}_n \cup \psi^{-1}\big((\alpha_n, L_n) \times S^1\big) \\ \Delta w_n = 0 & \text{on } \psi^{-1}\big((0, \alpha_n) \times S^1\big). \end{cases}$$

By the maximum principle, $\delta_n := \int_\Sigma w_n \, dg_n \leq \int_{\Sigma \setminus \mathcal{U}_n} dg_n$. Define $f_n = w_n - \delta_n$ so that $\int_\Sigma f_n = 0$ and f_n is admissible for the variational characterisation of $\lambda_1(g_n)$:

$$\lambda_1(g_n) \leq \frac{\int_\Sigma |\nabla f_n|^2 \, dg_n}{\int_\Sigma f_n^2 \, dg_n} \leq \frac{\int_D |\nabla w_n|^2 \, dg_{S^2}}{\int_\Sigma (w_n - \delta_n)^2 \, dg_n}.$$

The denominator satisfies

$$\begin{aligned} \int_\Sigma (w_n - \delta_n)^2 \, dg_n &\geq \int_{\mathcal{U}_n} (w_n^2 - 2\delta_n w_n + \delta_n^2) \, dg_n \\ &\geq \int_{\mathcal{U}_n} w_n^2 \, dg_n \\ &\geq \frac{1}{3} \int_{\mathcal{U}_n} dg_n. \end{aligned}$$

Whence,

$$\begin{aligned} \frac{\lambda_1(g_n)}{3} \int_{\mathcal{U}_n} dg_n &\leq \int_D |\nabla w_n|^2 \, dg_n \\ &\leq \int_{\mathcal{U}_n \cup \psi_n^{-1}(0,\alpha_n) \times S^1} |\nabla w_n|^2 \, dg_n + \int_{\psi_n^{-1}(\alpha_n, L_n) \times S^1} |\nabla w_n|^2 \, dg_n \\ &\leq \frac{8\pi}{3} + \int_{\psi_n^{-1}(\alpha_n, L_n)} |\nabla w_n|^2 \, dg_n. \end{aligned}$$

Proof of Theorem 1.1.3. The set $\psi^{-1}\big((0, \alpha_n) \times S^1\big)$ where w_n is harmonic is conformally equivalent to a cylinder of length $\alpha_n \geq \frac{L_n}{2}$ which becomes infinite as n goes to infinity. The proof is completed by choosing appropriate α_n as it was done in Lemma 3.6.1 and Lemma 4.5.2 and then applying Lemma 2.1.1 to bound the Dirichlet energy. $\square$

6. Acknowledgment

This work is a part of my PhD thesis at the Université de Montréal under the supervision of Marlène Frigon and Iosif Polterovich. I want to thank them for their support and encouragement. I would like to thank N. Nadirashvili for outlining the proof of Theorem 1.2.2 and 1.4.2 and for interesting discussions. I would also like to thank A. El Soufi, P. Guan, S. Ilias, M. Levitin, and M. Sodin for interesting and useful conversations. Finally, I want to thank my friends Baptiste Chantraine, Nicolas Beauchemin, Rémi Leclercq, and Jean-François Renaud for useful comments on an earlier version of this paper.

References

[1] M. Berger, P. Gauduchon and E. Mazet, *Le spectre d'une variété riemannienne*, Lecture Notes in Mathematics, Vol. **194**, Springer-Verlag, Berlin-New York, 1971.

[2] M. Berger, *Sur les premières valeurs propres des variétés riemanniennes*, Compositio Math., **26**, (1973), 129–149.

[3] S. S. Chern, M. do Carmo, S. Kobayashi, *Minimal submanifolds of a sphere with second fundamental form of constant length*, Functional Analysis and Related Fields (Proc. Conf. for M. Stone, Univ. Chicago, Chicago, Ill., 1968) pp. 59–75 Springer, New York, 1970.

[4] B. Colbois and J. Dodziuk, *Riemannian metrics with large λ_1*, Proc. Am. Math. Soc. **122**, (1994), 905–906.

[5] B. Colbois and A. El Soufi, *Extremal eigenvalues of the Laplacian in a conformal class of metrics: the 'conformal spectrum'*, Ann. Global Anal. Geom. **24**, (2003), no. 4, 337–349.

[6] A. El Soufi and S. Ilias, *Riemannian manifolds admitting isometric immersions by their first eigenfunctions*, Pacific J. Math. **195**, (2000), no. 1, 91–99.

[7] A. El Soufi, H. Giacomini and M. Jazar, *Greatest least eigenvalue of the laplacian on the klein bottle*, arXiv: math.MG/0506585 v1 29 Jun 2005.

[8] A. El Soufi, S. Ilias, *Immersions minimales, première valeur propre du laplacien et volume conforme*, Math. Ann. **275**, (1986), no. 2, 257–267.

[9] Hui Xia He and Zi Zhou Tang, *An isometric embedding of Möbius band with positive Gaussian curvature*, Acta Math. Sin. **20**, (2004), no. 6, 961–964.

[10] J. Hersch, *Quatre propriétés isopérimétriques des membranes sphériques homogènes*, C.R. Acad. Sc. Paris, Série A **270**, (1970), 1645–1648.

[11] A. Hurwitz, *Sur le probléme des isopérimètres*, Comptes rendus Acad. Sci. Paris, **132**, (1901), 401-403.

[12] D. Jakobson, M. Levitin, N. Nadirashvili, N. Nigam and I. Polterovich, *How large can the first eigenvalue be on a surface of genus two?*, Int. Math. Res. Not. (2005), no. 63, 3967–3985.

[13] D. Jakobson, N. Nadirashvili and I. Polterovich, *Extremal metric for the first eigenvalue on a Klein bottle*, Canad. J. Math. **58** (2006), no. 2, 381–400.

[14] H. B. Lawson, *Local rigidity theorems for minimal hypersurfaces*, Ann. of Math. (2) **89** 1969 187–197.

[15] S. Montiel, A. Ros, *Minimal immersions of surfaces by the first eigenfunctions and conformal area*, Invent. Math. **83** (1985), no. 1, 153–166.

[16] N. Nadirashvili, *Berger's isoperimetric problem and minimal immersions of surfaces*, GAFA **6**, (1996), 877–897.

[17] N. Nadirashvili and L. Friedlander, *A differential invariant related to the first eigenvalue of the Laplacian*, Internat. Math. Res. Notices **17**, (1999), 939–952.

[18] N. Nadirashvili, Private communication, 2005.

[19] P. Li and S.T. Yau, *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math. **69**, (1982), 269–291.

[20] R. Schoen, S.T. Yau, *Lectures on differential geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology, I. International Press, Cambridge, MA, 1994.

[21] P. Yang and S.T. Yau, *Eigenvalues of the Laplacian of compact Riemann surfaces and minimal submanifolds*, Ann. Scuola Norm. Sup. Pisa **7**, (1980), 55–63.

Département de Mathématiques et Statistique, Université de Montréal, C. P. 6128, Succ. Centre-ville, Montréal, Canada H3C 3J7

E-mail address: `girouard@dms.umontreal.ca`